

ORIGINAL ARTICLE

Transfusion Transmitted Infection Trends among Blood Donors in Tertiary Care Hospital from North India: A 7 Year Retrospective Study

Shailesh kumar Mishra¹, Shivani Kalhan² and Nand Kishore Singh²

¹Department of Pathology, ²Department of Community Medicine

^{1,2}Shaheed Hasan Khan Mewati (SHKM) Government Medical College Nalhar, Nuh (Mewat) Haryana-122107, India.

Abstract:

Background: Transmission of infection has always been a concern for blood bank professionals and for practising physicians who prescribe blood and blood components for their patients. The rate of Transfusion transmitted infections in donated blood is one of the main indicators of blood safety which has to be monitored precisely. This paper provides a thorough study of blood safety indicators. To study the trends of infections that can be transmitted by blood such as HIV, Hepatitis B Virus, Hepatitis C Virus, Syphilis and Malaria among blood donors at a tertiary care Government Medical College blood bank in north India in the last 7 years. **Material and Methods:** This study was conducted retrospectively over a period of 7 years from August 2013 to January 2020. A total of 29,262 blood donors who donated blood during this period were tested for HIV, HBV, HCV, Syphilis and Malaria. Comparative analysis was done to study the trend of transfusion transmitted infection among these blood donors. **Results:** : Out of 29,262 blood unit tested a total of 86 cases of HIV, 840 cases of HBV, 163 cases of HCV, 4 cases of syphilis and 13 cases of malaria were identified. Among the blood donors 0.294 % were reactive to HIV, 2.87% were reactive to HBsAg, 0.557 % were reactive to HCV. Whereas 0.0136 % were found to be positive for syphilis and 0.044 % were positive for malaria. **Conclusion:** Comparing various infections among repeat blood donor and first-time blood donor a retrospective data over a long period showed a trend in seroprevalance of HIV, HBsAg, HCV, Syphilis and malaria. The risk of transmission of these infections by blood transfusion remains high among first time donors. The results of the study reflect the prevalence of these infections in the apparently healthy population and warrant measures like effective donor retention and education policy and newer testing modality such as nucleic acid testing (NAT) to detect infection early and prevent residual risk of transmission.

Keywords: Blood Donor, HIV, Hepatitis B virus, Transfusion-transmitted infections (TTI), HBsAg, Prevalence.

Introduction:

Blood transfusion contributes to saving lives every day around the world, but it can also be a source of transmission of infection including HIV, HBV, HCV Syphilis and Malaria. Despite the necessity of blood components in certain medical circumstances, unsafe blood components can cause serious risks for the patients or even endanger their lives [1]. Although the performance of serological tests has been considerably improved in recent years; there remains a residual risk of transmission of viruses by blood transfusion. Blood presents a potential risk of infection to the receivers of transfusions. The rate of transfusion transmitted infections in donated blood plays a key role as the main indicator of blood safety, which has to be monitored precisely. The biggest challenge in this regard is transfusion transmitted infections (TTIs), the most important of which are human immune deficiency virus (HIV) as well as hepatitis types B (HBV) and C (HCV) viruses which create a large burden for the healthcare system [2,3]. Every blood transfusion carries potential risk of transmitting transfusion transmitted infections (TTI) to the recipient. Screening of blood donors first started in 1947 [4]. Today, India's blood transfusion program mandates the screening of HIV, hepatitis B virus (HBV), hepatitis C virus (HCV), malaria, and syphilis [5]. Estimated prevalence of HIV among adults in India is 0.2–0.3%; 6 and up to 40 million of the 350 million hepatitis B chronic carriers worldwide are in India [7].

Hepatitis B surface antigen (HBsAg) prevalence varies from 1% to 13%, with an average of 4.7% [8] and there are estimated 1-13 million HCV carriers in India [9]. Given the significance of these infections and the ease with which they are transmitted through blood components, the blood transfusion services follows a set of standard operating procedure (SOPs) to identify infected blood donations.

These procedures include two main activities: selecting safe donors and performing screening tests in order to identify TTIs [10]. The selection process include interviewing each blood donors by a qualified interviewer. In case of any risk factors or any history of risky behaviour found in blood donor, blood donation from these donors is refused [11,12]. In the next step, donated blood is screened for TTIs. These tests include screening for HCVAb, HBsAg and HIV (Ag/Ab) using the ELISA test, malaria by rapid card test method and syphilis by RPR method. These tests are performed according to the standard operating procedure of blood bank SHKM Government medical college guidelines [12] The screening method used to screen the donated blood needs to be improved over time based on new technologies of screening methods. For example, from ELISA to NAT can led to much more accurate and reliable test results and reduction in window period.

The present study was undertaken to study trends of transfusion transmitted infections such as HIV, Hepatitis B Virus, Hepatitis C Virus, Syphilis and Malaria among blood donors and compare the trend among first time blood donor and repeat blood donors.

Material and Methods:

This was a retrospective study conducted from August 2013 to January 2020 in the department of Blood Transfusion Shaheed Hasan Khan Mewati Government Medical College Nalhar, a tertiary care hospital in North India. All technical support was provided by the staff posted for the blood collection.

This study used the data registered in donor register that came for the blood donation either in blood bank or at voluntary blood donation camp. A total of 29,262 blood donor who fulfilled all the criteria for donation

were included in the study. The data includes the number of donors and the number of identified infections between August 2013 and January 2020. Blood donations and donor information collected for this study included years of collection, sex, age, status of donor (first time or repeated), collection site (fixed or mobile collection), and results (positive or negative) of serological tests for HIV, HBV, HCV, syphilis and malaria. A repeat donor was defined as a donor who had donated blood at least twice during the study period.

HIV, HBV, and HCV serological tests were performed according to an algorithm defined by the NACO. Initially, all donated blood was tested with fourth generation ELISA to detect anti-HIV1 and HIV2 antibodies and antigen P24 of HIV, surface-antigen HBs of HBV, and anti-HCV antibodies, respectively. Due to limited resources, positive samples were not confirmed by reference tests, such as Western blot for HIV or nucleic acid testing. Nevertheless, the algorithm applied by the NACO defined confirmatory strategies. For HIV, initial reactive samples were retested in duplicate using the same kit. Negative results in duplicate were considered negative. In cases of positive reaction, a parallel algorithm sampling, two rapid diagnostic tests (RDTs), a screening test and an HIV1 and HIV2 discriminant test was applied. Positive samples in both RDTs were considered positive. Discordant samples in both RDTs or between RDT and ELISA were considered indeterminate. For HBV and HCV, initial reactive samples were retested using HBsAg and SD Bioline HCV (Standard Diagnostics, Yongin, South Korea) respectively. Positive samples in both tests were considered positive, and discordant samples in both tests were considered indeterminate. Blood units detected positive or indeterminate for one TTI were discarded. Donors with indeterminate results were invited for a control test 3 months later.

If the sample tested positive for Syphilis and Malaria on rapid card test of different make then it was considered positive.

The statistical analysis was carried out using statistical package for social sciences (SPSS). Comparison for first time donors and repeat donors was done using

Mann-Whitney U-test. Proportions were compared using chi-square or Fisher’s exact test as applicable. All statistical tests were two sided and were performed at a significance level of 0.05.

Results:

A total of 29,262 units of blood was collected between August 2013 and January 2020 (66% were first time donors, and the remaining 34% had donated blood more than once, this ratio that has been almost constant in recent years) in SHKM GMC Nalhar (Table-1). Out of 29262 blood unit tested a total of 86 cases of HIV, 840 cases of HBV, 163 cases of HCV, 4 cases of syphilis and 13 cases of malaria were identified. Among the blood donors, 0.294 % was

reactive to HIV, 2.87% were reactive to HBsAg, 0.557 % were reactive to HCV. Whereas, 0.0136 % were found to be positive for syphilis and 0.044 % were positive for malaria (Table-2).

The period prevalence (7 years) of these infections was found to be 29, 287, 56, 1 and 4 cases per 10,000 donations for HIV, HBV, HCV, syphilis and malaria respectively. The largest numbers of infections were identified in the middle years of the study (2014 to 2017) whereas the number of infections was lowest in the year 2013. Among these infections, HBV had the highest prevalence (Table 2). Most number of transfusion transmitted infections were reported amongst blood donors with blood group “B”, this may be due to higher number of B blood group donors in the study (Table 3,4,5).

Table No. 1: Basic characteristics of the population under study

Parameter	N = 29262	%	P -Value
Sex			
Male	28,496	97.38	< 0.05
Female	766	2.62	
Age-Group (Years)			
< 20	2126	7.26	< 0.05
20 - 29	15,834	54.11	
30 - 39	7,891	26.96	
40 - 49	3134	10.71	
50 - 59	243	0.83	
60 - 65	34	0.11	
Donor Type			
First Time	19337	66.08	< 0.05
Repeat	9925	33.92	

Table No. 2: Year wise and total prevalence of transfusion Transmitted infections (TTIs)

TTIs ↓	YEAR →	2013	2014	2015	2016	2017	2018	2019	2020 (Jan)	Total (out of n=29, 262 Donation)	% Prevalence	P- Value
HIV (ELISA)		2	20	11	7	12	9	25	0	86	0.294	<.05
HBs Ag (ELISA)		20	95	94	106	167	160	189	9	840	2.87	
HCV (ELISA)		4	46	18	10	22	25	38	0	163	0.557	
Syphilis (RPR)		0	1	1	2	0	0	0	0	4	0.0136	
Malaria (Rapid card)		0	4	3	1	4	0	0	1	13	0.0444	

Table No. 3: Prevalence of TTI among different blood groups

Blood Group	Total Number of Blood unit	TTI Prevalence					Total no & %		P-Value
		HIV	HBs Ag	HCV	Syphilis	Malaria	Total	%	
A	6433	19	183	32	01	01	236	3.66	<.05
B	11465	44	439	79	02	09	573	4.99	
AB	2833	02	14	14	0	01	31	1.09	
O	8531	21	204	38	01	02	266	3.11	
Total	29262	86	840	163	4	13	1106	3.78	

Table No. 4: Year-wise Prevalence of different blood groups

Blood Group	2013	2014	2015	2016	2017	2018	2019	2020 (January)	Total	% of Total	P-Value
A	217	601	807	916	1103	1327	1364	98	6433	21.98	<.05
B	358	1098	1375	1556	2103	2215	2584	176	11465	39.19	
AB	85	241	408	383	509	581	582	44	2833	9.67	
O	293	836	1050	1162	1545	1719	1794	132	8531	29.16	
Total	953	2776	3640	4017	5260	5842	6324	450	29262	100	

Table No. 5: Distribution of Rh (positive) and Rh (negative)

Blood Group	Total Blood unit collected	%	P-Value
Rh (positive)	26903	91.94	<.05
Rh (negative)	2359	8.06	
Total	29262	100%	

Table No. 6: Comparative % prevalence among different region of India

Region	Place	HIV (%)	HBs Ag (%)	HCV (%)	Syphilis (%)	Malaria (%)	P-Value	Author and Year
Present Study	South Haryana	0.294	2.87	0.556	0.0136	0.0444	< 0.05	Shailesh et al. 2013-2020
North India	Ludhiana Punjab	0.08	0.66	1.09	0.85	--	< 0.05	Gupta et al. 2001-2003 [17]
	Etawah UP	0.19	2.63	0.34	--	--		Dayal S et al. 2006-2012 [6]
	Lucknow UP	0.23	1.96	0.85	0.01	--		Chandra et al. 2001-2006 [27]
	Bikaner Rajasthan	0.10	1.60	0.18	0.89	0.04		Devraj et al. 2001-2003 [28]
Central India	Bhanpura MP	0.51	2.90	0.57	0.23	--	< 0.05	Sawke et al. 2006-2008 [29]
South India	Vikrabad AP	0.39	1.41	0.84	0.08	--		Bhawani et al. 2004-2009 [30]

Discussion:

Blood transfusion services not only screen the blood donor but also give a clue about the rate of prevalence of TTI in asymptomatic healthy young adults. It can be considered as a reliable tool for statistical estimation of these infections in the general population as discussed by Gharehbaghian, Attaullah, et al. [13,14]. When basic characteristics of the population under study was analysed, it was found that maximum transfusion transmitted infections (TTIs) were reported among age group of 20-29 years. It may be because of higher number of blood donations by individuals in this age group (Table 1). The prevalence rate may be under or overestimate due to their different gender, age, area of residence, and other characteristics [15,16]. Blood transfusion is an important preventable modality of spread of TTI. Accurate estimation of risk of TTIs is essential for monitoring of blood safety and efficacy of currently employed screening programs as discussed by NACO guidelines [17].

The outcome of the present study can suggest to formulate strategies to identify, recruit, and retain VBD (Voluntary Blood Donor) from low risk population, as blood donors with high risk group such as donor with history of jaundice, injectable drug abuse, and multiple sexual partners can be screened out by donor questionnaire. It has been found that prevalence of various TTI varies among different geographical regions [14].

The results indicated that the prevalence of HBV (287 cases per 10,000 donations) was higher than that of HCV (56 cases per 10,000 donations) in the study period (7 years) while the prevalence of HIV (29 cases per 10,000 donations) was comparable to that in blood donors in most other countries. 31 Other studies in India (HBV=1.77%, HIV=0.14% and HCV=0.04%), Colombia (HBV=0.12%, HCV=0.45% and HIV=0.12%), Italy (HBV=0.01%) and Turkey (HBV=4.19% and HCV=0.38%) showed a higher prevalence of TTIs among blood donors compared to this study [18-21]. A systematic review of studies on the prevalence of HBV among the WHO member countries showed that the prevalence of HBV in Italy (2.52%), Colombia (2.3%) and Turkey (4%) is

comparable to our present study (2.88%) [22]. The prevalence of TTIs among blood donors in this study showed a decreasing trend between the years 2013 and 2020. The results of other studies on the prevalence trends of TTIs among blood donors of other study are similar to the results of our study and show a decreasing trend of TTIs [1,23]. This study showed that highest prevalence of TTIs was observed in the first-time donors which is similar to Iranian study [24]. It appears that as years go by, a larger ratio of donors is made up from repeat and regular blood donors. These people received more education compared to first-time blood donors, which can explain a lower prevalence of TTIs among repeat donors. When various other studies from different parts of India were compared, it was found that percent prevalence of HIV was similar to other studies, though some studies reported less prevalence specially study from Maharashtra and Ludhiana, Punjab. Percent prevalence of HBsAg were similar to the study done at Bhanpura MP (2.9%). All other studies had lower prevalence compared to present study. Study done in West Bengal and Etawah UP showed less prevalence of HCV than present study and all other studies showed higher prevalence for HCV compared to present study. Study done at Ludhiana Punjab (0.85%), Bikaner Rajasthan (0.89%), Karnataka (1.6%), West Bengal (0.72%), Bhanpura MP (0.23%) showed higher percent prevalence for syphilis compared to present study whereas study done at Lucknow and Vikarabad AP showed similar result for percent prevalence for HCV compared to present study. This study showed highest prevalence of malaria compared to other study (Table 6).

Conclusion:

We have observed decreasing trend of HIV, HBsAg, HCV, syphilis and malaria which is significant in repeat blood donor as compared to first time blood donors. Voluntary donations are safer as compared to replacement ones and should be encouraged. Efforts should be made to increase the number of voluntary donors with a target of 100% and reduce replacement donations to a minimum. Transmission of TTIs during serologically negative window period is still a threat to

blood safety. Therefore, strict selection of blood donors and comprehensive screening of donor's blood using standard and latest technique such as NAT are highly recommended to ensure the safety of blood for recipient. This trend suggests that for prevention of

TTIs by healthy VBD recruitment and safe VBD retention strategies should be strengthened at all levels in blood transfusion service so that first time blood donors may turn into repeat blood donor.

Conflict of Interest - Nil

Sources of Support - Nil

References

- Mohammadali F, Pourfathollah AA. Changes in frequency of HBV, HCV, HIV and syphilis infections among blood donors in Tehran province 2005–2011. *Archives of Iranian Medicine* 2014;17(9):613–620.
- Flichman DM, Blejer JL, Livellara BI, et al. Prevalence and trends of markers of hepatitis B virus, hepatitis C virus and human immunodeficiency virus in Argentine blood donors. *BMC Infectious Diseases* 2014;14(1):218.
- Mohsenizadeh M, Mollaei HR, Ghaziizadeh M. Seroepidemiological study of hepatitis B, C and HIV among blood donors in Kerman. *Asian Pacific Journal of Cancer Prevention* 2017;18(12):3267.
- Available from: <http://www.redcrossblood.org/learn>
- National AIDS Control Organization (NACO, India). Standards for Blood Banks & Blood Transfusion Services. New Delhi: NACO, Ministry of Health and Family Welfare, Government of India; 2007:11.
- Available from: <http://www.unaids.org/en/data-analysis/datatools/aidsinfo/>.
- Datta S. An overview of molecular epidemiology of hepatitis B virus (HBV) in India. *Virology Journal* 2008;5:156.
- Sachan D, Varghese J, Joseph J, Srinivasan V, Jayanthi V, Rela M. HBsAg positive patient characteristics in hospital and blood donation camps. *ISRN Hematology* 2013 Sept 13; 2013: 675191.
- Narahari S, Juwle A, Basak S, Saranath D. Prevalence and geographic distribution of hepatitis C virus genotypes in Indian patient cohort. *Infection, Genetics and Evolution* 2009;9:643–645.
- Saber HR, Tabatabaee SM, Abasian A, et al. Incidence and residual risk of HIV, HBV and HCV infections among blood donors in Tehran. *Indian Journal of Hematology and Blood Transfusion* 2017;33(3):412–416.
- Tessema B, Yismaw G, Kassu A, et al. Seroprevalence of HIV, HBV, HCV and syphilis infections among blood donors at Gondar University Teaching Hospital, Northwest Ethiopia: declining trends over a period of five years. *BMC Infectious Diseases* 2010;10(1):1–7.
- Sajjadi SM, Pourfathollah AA, Mohammadi S, Nouri B, Hassanzadeh R, Rad F. The prevalence and trends of hepatitis B, hepatitis C, and HIV among voluntary blood donors in Kohgiluyeh and Boyer-Ahmad transfusion center, Southwestern Iran. *Iranian Journal of Public Health* 2018;47(7):944.
- Gharehbaghian A. An estimate of transfusion-transmitted infection prevalence in general populations. *Hepatitis Monthly* 2011;11(12):1002–1003.
- Attaullah S, Khan S, Khan J. Trend of transfusion transmitted infections frequency in blood donors: Provide a road map for its prevention and control. *Journal of Translational Medicine* 2012;10:20.
- Ali SA, Donahue RM, Qureshi H, Vermund SH. Hepatitis B and hepatitis C in Pakistan: Prevalence and risk factors. *International Journal of Infectious Diseases* 2009;13:9–19.
- Jafri W, Jafri N, Yakoob J, Islam M, Tirmizi SF, Jafar T, et al. Hepatitis B and C: Prevalence and risk factors associated with seropositivity among children in Karachi, Pakistan. *BMC Infectious Diseases* 2006;6:101.
- Ministry of health and family welfare government of India New Delhi. Voluntary Blood Donation Programme – An Operational Guideline. New Delhi: National AIDS Control Organization, Government of India; 2007.
- Omez LA, Penuela O, Higuera F. Prevalence of antibodies against transfusion-transmissible infections (TTI) in blood donors from the Colombian eastern region. *Clinical Laboratory* 2014;60(5):869–871.
- Gurol E, Saban C, Oral O, Cigdem A, Armagan A. Trends in hepatitis B and hepatitis C virus among blood donors over 16 years in Turkey. *European Journal of Epidemiology* 2006;21(4):299–305.
- Yadav B, Varma A, Singh P, Kumar R, Bandi P. Seroprevalence of Transfusion-Transmitted Infections (TTIs) in blood donors: a study from Central India. *International Journal of Medical Science and Public Health* 2016;5(6):1–5.
- Romanò L, Velati C, Cambiè G, et al. Hepatitis B virus infection among first-time blood donors in Italy: prevalence and correlates between serological patterns and occult infection. *Blood Transfusion* 2013;11(2):281.
- Schweitzer A, Horn J, Mikolajczyk RT, Krause G, Ott JJ. Estimations of worldwide prevalence of chronic hepatitis B

- virus infection: a systematic review of data published between 1965 and 2013. *Lancet* 2015; 386 (10003):1546–1555.
23. AminiKafi-abad S, Rezvan H, Abolghasemi H, Talebian A. Prevalence and trends of human immunodeficiency virus, hepatitis B virus, and hepatitis C virus among blood donors in Iran, 2004 through 2007. *Transfusion* 2009;49(10):2214–2220.
 24. Mahdavian F, Saremi S, Maghsoudlu M, Pourfathollah AA. Prevalence of blood transmitted viral infections in regular and non-regular donors of Arak Blood Centre. *Sci JIBTO* 2006;2 (7):343–351.
 25. Gupta N, Kumar V, Kaur A. Seroprevalence of HIV, HBV, HCV and syphilis in voluntary blood donors. *Indian Journal of Medical Sciences* 2004;58:255-257.
 26. Dayal S, Omar BJ, Sipai HH, Agarwal SK, Chandra S, Chaturvedi V. Prevalence of transfusion transmitted viral diseases among blood donors in a rural setup of North India from 2006–2011. *International Journal of Science, Technology and Management* 2011;2:97-99.
 27. Chandra T, Kumar A, Gupta A. Prevalence of transfusion transmitted infections in blood donors: An Indian experience. *Transfusion* 2009;49:2214-2220.
 28. Arya D, Mahawar NL, Pachaury R, Bharati A, Sharma L, Kumar H, Singh VB. Seroprevalence of transfusion transmitted infections among blood donors at a tertiary care Hospital Blood Bank in North India. *Indian Journal of Health Sciences* 2016;9(1):77-81
 29. Sawke N, Sawke GK, Chawla S. Seroprevalence of common transfusion –Transmitted infections among blood donors at Bhanpur, Bhopal, M.P. from 2006–2008. *Peoples Journal of Science and Research* 2010;3:5-7.
 30. Bhawani Y, Rao PR, Sudhakar V. Seroprevalence of transfusion transmissible infections among blood donors in a tertiary care hospital of Andhra Pradesh from 2004–2009. *Biology and Medicine* 2010;2:45-48.
 31. Heyredin I, Mengistie B and Weldegebreal F. Seroprevalence of transfusion-transmittable infections and associated factors among blood donors in Eastern Ethiopia: an Institutional-based cross-sectional study. *SAGE Open Medicine* 2019; 7: 1–8.

Address for correspondence: Dr. Shailesh Kumar Mishra, Assistant Professor of Pathology, Shaheed Hassan Khan Mewati Government Medical College, Nalhar, Nuh, Haryana-122107 India. Email: dr.shaileshmishra@gmail.com, Mobile: +91 7982708046.

How to cite this article: Shailesh kumar Mishra, Shivani Kalhan and Nand Kishore Singh. Transfusion Transmitted Infection Trends among Blood Donors in Tertiary Care Hospital from North India: A 7 Year Retrospective Study. *Walawalkar International Medical Journal* 2021; 8(1):52-58. <http://www.wimjournal.com>

Received date: 23/01/2021

Revised date: 12/06/2021

Accepted date: 14/06/2021